



**Orthopaedic Associates of Zanesville**  
**Post-Op Instructions Trigger Finger Release**  
**Dr. Brautigam**

Please note that these instructions are general guidelines to be followed; however, any written or verbal instructions provided by your provider supersede the instructions below and should be followed.

**WOUND CARE**

- It is common to have leakage of fluid and blood into or around the dressing the first 48 hours after surgery. Do not remove the dressing.
- You are permitted to shower on post-op day 3. Please do not remove the dressing.

**SWELLING**

- Swelling is normal and expected. The dressing and any wrappings are placed on very loosely to allow for swelling. Sometimes the swelling can be painful due to the wrappings, if so please call our office for instructions. Usually just loosening the wrappings will make a difference. Continue to elevate the leg above the level of the heart to help with the swelling.

**MEDICATION**

- You will typically be prescribed one of the following medication to be used after surgery at home for pain. Please take these how they are prescribed on the actual prescription:  
Percocet (oxycodone / acetaminophen): A strong narcotic pain reliever. Do not exceed 10 tablets in a 24 hour period.

OR

Over the counter Tylenol 500-1000mg every 6 hours as needed for pain and may alternating with ibuprofen 600-500mg every 6 hours as needed for pain.

- Keep in mind that all narcotics will cause drowsiness and should not be used with alcohol or if driving. Taking this medication on an empty stomach can result in an upset stomach. It is common to feel the most discomfort when lying down to sleep at night, so you may find it beneficial to take the medication at bedtime.
- If unusual side effects occur, please discontinue the medication and contact our office.

**EMERGENCIES**

- With any surgery it is important to be aware of any signs of infection, which can include: unusual looking or smelling drainage (green/yellow), increased redness and tenderness around the incision site, or fever (i.e. 101 degrees or above) and chills. Also, be aware of excessive bleeding or continued numbness and tingling. If you experience any of these symptoms, please call our office. There is a doctor on call 24 hours a day.

**FOLLOW-UP CARE/QUESTIONS**

- If you do not already have a post-operative appointment scheduled, please contact our office at 740-454- 3273 to schedule.
- Typically, the first post-operative appointment following surgery is 10-14 days following surgery. If you have any further questions please contact us during office hours.